



IGH Distal Plus Probe


REF: LPH 516-A/LPH 516-A50

Analyte Specific Reagent: Analytical and performance characteristics are not established.

Fluorescence $in\ situ$ hybridisation (FISH) is a technique that allows the visualisation of DNA sequences upon chromosomes. The technique uses DNA probes that hybridise to entire chromosomes or single unique sequences, and serves as a powerful adjunct to classic cytogenetics. Recent developments have meant that this valuable technique can now be applied as an essential tool in prenatal, haematological and pathological chromosomal analysis. Target DNA, after fixation and denaturation, is available for annealing to a similarly denatured, fluorescently labelled DNA probe, which has a complementary sequence. Following hybridisation, unbound and non-specifically bound DNA probe is removed and the DNA is counterstained for visualisation. Fluorescence microscopy then allows the visualisation of the hybridised probe on the target material.

Probe Specification

IGHV, 14q32.33, Green



The IGH Distal Plus product consists of a 324kb probe and a 289kb probe within the Variable segment of the IGH region

Materials Provided

Probe: 50µl per vial or 250µl per vial

Probe concentration: Amount of green probe 4.80-7.18ng/µl

The probe is provided in hybridisation solution (Formamide; Dextran Sulphate; SSC) and is ready to use.

Warnings and Precautions

- For professional use only.
- Wear gloves when handling DNA probes.
- Probe contains formamide, which is a teratogen; do not breathe fumes or allow skin contact. Wear gloves, a lab coat, and handle in a fume hood. Upon disposal, flush with a large volume of water.
- Dispose of all hazardous materials according to your institution's guidelines for hazardous waste disposal.
- Operators must be capable of visually distinguishing between red, blue and 5 areen.

Storage and Handling

The kit should be stored between -25°C to -15°C in a freezer until the expiry date indicated on the kit label. Store the probe vial in the dark. Ensure that exposure of the probe to laboratory lights is limited at all times.

Known Cross-Reactivity

The green IGH probe may show cross-reactivity to 15q11.2 and 16p11.2.

Additional Information

For additional product information please contact the CytoCell Technical Support Department.

T: +44 (0)1223 294048

E: techsupport@cytocell.com

W: www.ogt.com

Labelling according to Regulation (EC) No. 1272/2008 [CLP]

Hazard pictograms (CLP):





GHS07

GHS08

Signal word (CLP): Danger

Hazardous ingredients: Formamide <100%

Hazard statements (CLP):

H315 - Causes skin irritation

H319 - Causes serious eye irritation

H360 - May damage fertility or the unborn child

Precautionary statements (CLP):

P202 - Do not handle until all safety precautions have been read and understood P280 - Wear eye protection, protective clothing, protective gloves P302+P352 - IF ON SKIN: Wash with plenty of soap and water P305+P351+P338 - IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing P308+P313 – IF exposed or concerned: Get medical advice/attention P362+P364 - Take off contaminated clothing and was it before reuse P501 - Dispose of contents/container to hazardous or special waste collection point, in accordance with local, regional, national and/or international regulation

Refer to Safety Data Sheet for more information.

Patents and Trademarks

CytoCell is a registered trademark of Cytocell Ltd.



Cytocell Ltd.

Oxford Gene Technology, 418 Cambridge Science Park, Milton Road, Cambridge, CB4 0PZ, UK T: +44(0)1223 294048 F: +44(0)1223 294986 E: probes @cytocell.com W: www.ogt.com